

Cattle Dewormer Comparison Chart

	Active Ingredient	Lungworm	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Sucking Lice	Biting Lice	Mange Mites	Horn Flies	Grubs	Beef	Lactating Dairy	Non-Lactating Dairy	Calves	Indications	Dosage	Slaughter Withdrawal	Route of Administration
Injectable																								
Agri-Mectin Cattle/Swine Injection	ivermectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ
Bimectin Cattle/Swine Injection	ivermectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ
Bimectin Plus Injection	ivermectin & clorsulon	•	•				•		•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	21 days	SQ
Cydetin Cattle Injectable	moxidectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	21 days	SQ
Dectomax Cattle/Swine Injection	doramectin	•	•						•	•	•	•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ, IM
Ivermectin Cattle/Swine Injection	ivermectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ
Ivermectin Plus Injection	ivermectin & clorsulon	•	•				•		•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	21 days	SQ
Ivomec Cattle/Swine Injection	ivermectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ
Ivomec Plus Cattle Injection	ivermectin & clorsulon	•	•				•		•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	49 days	SQ
LongRange Injectable	eprinomectin	•	•	•	•	•			•					•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	48 days	SQ
Noromectin Cattle/Swine Injection	ivermectin	•	•						•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	35 days	SQ
Noromectin Plus Cattle Injection	ivermectin & clorsulon	•	•				•		•			•		•		•	•	•	•		Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 110 lbs	49 days	SQ

Cattle Dewormer Comparison Chart

	Active Ingredient	Lungworm	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Sucking Lice	Biting Lice	Mange Mites	Horn Flies	Grubs	Beef	Lactating Dairy	Non-Lactating Dairy	Calves	Indications	Dosage	Slaughter Withdrawal	Route of Administration
Drench/Oral																								
DuraFend Type B & C	fenbendazole	•	•	•	•	•											•	•	•	•	Do not use in calves to be processed for veal	1 lb per 1000 lbs	13 days	Pellets
DuraFend Type C	fenbendazole	•	•	•	•	•			•								•	•	•	•	Do not use in calves to be processed for veal	1 lb per 1000 lbs	13 days	Pellets
Levamed Soluble Drench	levamisole HCl	•	•	•	•	•			•								•		•	•	Do not use in female dairy cattle over 20 months of age	2 ml per 100 lbs	2 days	Drench
Prohibit Soluble Drench	levamisole HCl	•	•	•	•	•			•								•		•	•	Do not use in female dairy cattle over 20 months of age	2 ml per 100 lbs	2 days	Drench
Safe-Guard 0.5% Pellets	fenbendazole	•	•	•	•	•			•								•	•	•	•		0.5 lbs per 500 lbs	13 days	Pellets
Safe-Guard 32SG Mineral Type C Medicated Feed	fenbendazole	•	•	•	•	•			•								•					1.6 oz per 100 lbs	13 days	Free-Choice
Safe-Guard Cattle Block	fenbendazole	•	•	•	•	•			•								•		•	•	Do not use in female dairy cattle over 20 months of age	2.27 mg per lb	16 days	Free-Choice
Safe-Guard Paste	fenbendazole	•	•	•	•	•			•								•	•	•	•	Do not use in calves to be processed for veal	5 gm per 220 lbs	8 days	Oral
Safe-Guard Suspension	fenbendazole	•	•	•	•	•			•								•	•	•	•	Do not use in calves to be processed for veal	2.3 ml per 100 lbs	8 days	Drench
Synanthic Suspension	oxfendazole	•	•	•		•		•	•								•		•	•	Do not use in female dairy cattle over 20 months of age	1 ml per 110 lbs	7 days	Drench
Valbazen Suspension	albendazole	•	•	•	•	•	•	•									•		•	•	Do not use in female dairy cattle over 20 months of age	4 ml per 100 lbs	27 days	Drench

Cattle Dewormer Comparison Chart

	Active Ingredient	Lungworm	Stomach Worms	Intestinal Worms	Bankrupt Worms	Nodular Worms	Liver Flukes	Tapeworms	Roundworms	Kidney Worms	Eyeworms	Sucking Lice	Biting Lice	Mange Mites	Horn Flies	Grubs	Beef	Lactating Dairy	Non-Lactating Dairy	Calves	Indications	Dosage	Slaughter Withdrawal	Route of Administration	
Pour-On																									
Bimectin Cattle Pour-On	ivermectin	•	•						•			•	•	•	•	•	•		•	•	•	Do not use in female dairy cattle over 20 months of age or veal calves	1 ml per 22 lbs	48 days	Topical
Cydectin Cattle Pour-On	moxidectin	•	•						•			•	•	•	•	•	•	•	•	•	•	Do not use in calves to be processed for veal	5 ml per 110 lbs	0 days	Topical
Dectomax Cattle Pour-On	doramectin	•	•	•		•			•			•	•	•	•	•	•	•	•	•	•	Do not use in female dairy cattle over 20 months of age or veal calves	5 ml per 110 lbs	45 days	Topical
Eprinex Cattle Pour-On	eprinomectin	•	•						•			•	•	•	•	•	•	•	•	•	•	Do not use in calves to be processed for veal	5 ml per 110 lbs	0 days	Topical
Eprizero Cattle Wormer Pour-On	eprinomectin	•	•						•			•	•	•	•	•	•	•	•	•	•	Do not use in calves to be processed for veal	5 ml per 110 lbs	0 days	Topical
Ivermectin Pour-On for Cattle	ivermectin	•	•						•			•	•	•	•	•	•		•	•	•	Do not use in female dairy cattle over 20 months of age or veal calves	5 ml per 110 lbs	48 days	Topical
Ivomec Cattle Pour-On	ivermectin	•	•						•			•	•	•	•	•	•		•	•	•	Do not use in female dairy cattle over 20 months of age or veal calves	5 ml per 110 lbs	48 days	Topical